DYSTOCIA DUE TO ABNORMAL CALF IN A MURRAH BUFFALO: A CASE REPORT

Parthasarathi Balasamudram Chandrasekhar^{1,*}, Dayanidhi Jena² and Pavankumar Gajula³

ABSTRACT

The fetal abnormalities are one of the protagonist causes for causing parturient related issues ultimately resulting in dystocias. Several congenital deformities observed in cattle and buffaloes and among those hydrocephalus, bulldog calf conditions are common. In this case report the handling, obstetrical correction, its post-delivery care and therapeutic medications are discussed.

Keywords:Bubalusbubalis,buffalo,hydrocephalus, congenital defect, dystocia

INTRODUCTION

There are several developmental abnormalities have been reported in domestic animals from time to time (Christopher, 1971; Jena *et al.*, 2016) leading to the dystocia condition and other peri and post parturient conditions and hydrocephalus anomaly is one amongst them. Hydrocephalus anomaly occurs due to accumulation of fluids causes dilatation of subarachnoid space and ventricular system (Noakes et al., 2009). During the foetal development, any obstruction in cerebrospinal fluid passage causes excessive accumulation in arachnoid space leads to over swelling of cranial cavity (Salunke et al., 2001). This condition also caused due to environmental nutritional and genetic factors (Kalman, 1989). This condition most commonly observed in cattle (Jana and Ghosh, 2005; Purohit et al., 2006). Calves with pronounced cranial enlargement usually die within 48 h but less severely affected calves may survive for several weeks or longer (Leech et al., 1978). The Bulldog condition common in foetuses showing hydrocephalus condition. The bulldog appearance due to shortened upper jaw, compressed skull and nose divided by furrows, as per Noakes (loc. cit.), this condition is uncommon in buffaloes.

CASE HISTORY

A full term pregnant Murrah buffalo in its first parity presented to Veterinary hospital with history of labour pain for one 28 h. Ten to sixteen hours before water bag had ruptured and one foetal

¹Division of Parasitology, Indian Veterinary Research Institute, Uttar Pradesh, India, *E-mail: parthb763@gmail.com

²Division of Animal Reproduction, Indian Veterinary Research Institute, Uttar Pradesh, India ³Veterinary Assistant Surgeon, Telangana, India leg was protruded out of the vulva. The animal was exhausted due to severe straining. On per vaginal examination the foetus positioned in anterior longitudinal presentation, dorso-sacral position with one fore-limb protruded from vulva and other leg was flexed from shoulder joint. The head of the foetus felt deformed, foetal movements and other reflexes were absent.

TREATMENT AND DISCUSSION

The animal was prepared for the manual correction of the postural defect. After sufficient lubrication, the flexed limb was brought to the passage in the normal posture manually. With using of obstetrical chains limbs were snared and traction was applied to deliver the fetus. A dead male foetus was delivered. The calf showed typical bulldog facial features (Figure 1) of shortened upper jaw, nose divided by furrows, protruding tongue, prolonged lower jaw and shortened upper jaw so this foetus was diagnosed as "bulldog calf" as described by Noakes, 2009. Moreover, the fetus was also affected with hydrocephalus condition as shown in Figure 1. Following delivery, dam treated with ecbolic agents to hasten the expulsion of placenta, antihistaminic preparation, a dose of antibiotics and anti inflammatory drugs for treatment and prophylactic measure.

A simple, autosomal recessive defects with some modifiers may cause the bull dog condition in animals (Roberts, 1986). Dystocia due to bull dog calf condition is uncommon in domestic animals but very few cases have been reported in cattle (Roberts, 1986; Harper *et al.*, 1998) and in buffaloes (Christopher, 2000); Shukla *et al.*, 2007).



Figure 1. Bulldog buffalo calf with hydrocephalus.

REFERENCES

- Christopher, K.J. 1971. Some developmental abnormalities in domestic mammals. J. Anat. Soc. India, 20: 33-35.
- Christopher, K.J. 2000. Some developmental abnormalities in buffalo. *Buffalo Bull.*, **19**(3): 64-66. Available on: https://ibic.lib. ku.ac.th/e-Bulletin/IBBU200000015.pdf
- Christopher, K.J. and K.B. Singh. 1997. Further observations on the developmental abnormalities of buffaloes. *Buffalo Bull.*, 16(1): 11-14. Available on: https://ibic.lib.ku.ac.th/e-Bulletin/IBBU200000002.pdf
- Harper, P., M.R. Latter, F.W. Nicholas, R.W. Cook and P.A. Gill. 1998. Chondrodysplasia in Australian dexter cattle. *Aust. Vet. J.*, 76(3): 199-202. DOI: 10.1111/j.1751-0813.1998. tb10129.x.
- Jana, D. and M. Ghosh. 2005. Hydrocephalus in a calf: A case report. *The North East Veterinarian*, **14**(4): 11.
- Jena, D., B.K. Patra, S. Das and D.N. Mohanty. 2016. Acephalic monster- A case report. *Indian Vet. J.*, **93**(2): 85-86.
- Kalman, T.S. 1989. Congenital Malformations in Laboratory and Farm Animals. Academic Press. INC San Diego, California, USA.
- Leech, R.W., C.N. Hauges and L.A. Christoferson.
 1978. Hydrocephalus, congenital hydrocephalus. Aminal model: Bovine hydrocephalus, congenital internal hydrocephalus, aqueductal stenosis. *Amer. J. Pathol.*, **92**(2): 567-570. Available on: https://www.ncbi.nlm.nih.gov/pmc/articles/ PMC2018286/pdf/amjpatho100736-0251. pdf
- Noakes, D.E., T.J. Parkinson and G.C.W. England. 2009. Veterinary Reproduction and

Obstetrics, 9th ed. Saunders London, UK. p. 950.

- Purohit, G.N., M. Gaur and A. Sharma. 2006. Hydrocephalus in a cow: A case report. Indian J. Anim. Reprod., 27: 98-99.
- Roberts, S.J. 1986. *Veterinary Obstetrics and Genital Diseases*, 2nd ed. CBS Publishers and Distributors, Delhi, India. p. 51.
- Salunke, S.P., M.B. Amle and P.C. Zambre. 2001. Dystocia due to hydrocephalus in Pantharpui buffalo. *Indian J. Anim. Reprod.*, **22**: 96.
- Shukla, S.P., S.P. Nema, A.K. Pandey and U.K. Garg. 2007. Dystocia due to bull dog calf in a she buffalo. *Buffalo Bull.*, 26(3): 104-105. Available on: https://ibic.lib.ku.ac.th/e-Bulletin/IBBU200700021.pdf