

SURVEY OF EXISTING RURAL FEEDING PRACTICES AND
NUTRIENT AVAILABILITY TO LACTATING BUFFALOES
IN INDORE DISTRICT OF MADHYA PRADESH

Anil Kumar Yadav*, Rvindra Kumar Jain, Dinesh Thakur and Mukesh Kumar Mehta

ABSTRACT

In Indore district of Madhya Pradesh, farmers are following traditional feeding practices. Dairy animals are mostly fed on straw based ration. Due to non-availability of balanced ration to dairy animals, productive performance is not up to the level of the satisfaction. Keeping this in view the survey of existing feeding practices in lactating buffaloes was done in the 10 villages around College of Veterinary Science and Animal Husbandry, Mhow. Eight to twelve farmers from each village were selected randomly and information about herd strength, available feedstuffs, use of mineral-vitamin supplements, daily feed offered and residue left were collected in a questionnaire. Milk yield of individual animal was also recorded. Quantities of feed offered/residue left and milk yield were verified randomly by weighing through portable digital balance. Milk samples were collected and analysed for fat content, randomly. Body weight of animals was determined from body measurements. The representative sample of each feed ingredient was collected and analyzed for proximate principles, Ca and P. The carotene was calculated from reported values of feedstuffs. Then nutrient (DCP, TDN, Ca, P and Carotene) supply was calculated

and compared with standard requirements to work out nutrient excess/deficit. It has been observed that five types of feeding patterns were followed by farmers, out of 104 dairy farmers visited having 347 buffaloes. Animals were fed in stalls two times a day. The available feedstuffs were straws (wheat/gram/masoor/soybean) uncultivated natural local grass, MP chari, maize fodder, cotton seed cake (un-decorticated), wheat bran and compounded feed. Majority of farmers were feeding mixture of straws (wheat straw + gram straw) + local grasses + cotton seed cake + wheat bran. Very few farmers were supplementing mineral mixture and salt. No farmer is supplementing vitamins. Studies on nutrient availability in different feeding patterns indicated that DCP was excess (4.91 to 28.52%), while TDN (3.28 to 10.93%), phosphorous (18.17 to 43.73%) and carotene (17.90 to 57.54%) were deficient in all five feeding patterns. Calcium was deficient in pattern I (4.16%), II (54.83%) and III (17.16%) and excess in feeding pattern IV (8.03%) and pattern V (65.66%). Ca : P ratios in rations of buffaloes under different feeding pattern ranged between 1.25:1 to 3.55:1 against optimum ratio 1.56 : 1. It was concluded that DCP was moderately high and TDN was marginally deficient while phosphorous and carotene were moderate to highly deficient in the rations of lactating buffaloes of the

region. Calcium was deficient in feeding pattern I, II and III and adequate in pattern IV while excess in pattern V.

Keywords: *Bubalus bubalis*, buffaloes, feeding practices, nutrient status, lactating buffaloes, Madhya Pradesh, India

INTRODUCTION

Most of the dairy animals in India are fed on agriculture by-products and low quality crop residues, which have low nutritive value and digestibility. The shortage of feed resources coupled with their poor nutritive value is the main reason of low productivity of dairy animals. On other hand, Imbalanced feeding leads to excess feeding of some nutrients while others remain deficient. This not only reduces milk production and increases costs per kg milk, but also affects various physiological functions including long term animal health, fertility and productivity (Garg *et al.*, 2000).

In Indore district of Madhya Pradesh, farmers are following traditional feeding practices. Dairy buffaloes are mostly fed on straw based ration with insufficient quantity of imbalanced concentrate mixture without mineral and/or vitamin supplementation. Seasonal green fodders like maize, MP chari, local green grasses during kharib and berseem and lucerne during rabi were offered in limited quantity depending upon the availability (Tewari *et al.*, 2012). Under such feeding situation, the availability of various nutrients which are having the importance in health and production may greatly reduce to animals. Keeping in view the above facts, present study was conducted to observe the existing feeding practices

and nutrient availability in lactating buffaloes under field conditions which will be useful to suggest the appropriate nutrient supplementation for enhancing productivity and profit to farmers.

MATERIALS AND METHODS

Survey of existing feeding practices in lactating buffaloes was done in the 10 villages (Bhaislai, Sonwai, Borkhedi, Panda, Navda, Harsola, Santer, Kavti, Rangwasa and Umriya) around College of Veterinary Science and Animal Husbandry, Mhow. Eight to twelve farmers from each village were selected randomly and information about herd strength, available feedstuffs, use of mineral-vitamin supplements, daily feed offered and residue left were collected in a questionnaire. Milk yield of individual animal was also recorded. Quantities of feed offered/ residue left and milk yield were verified randomly by weighing through portable digital balance. Milk samples were collected and analyzed for fat content, randomly by using automatic fat tester. Body weight of animals was determined from body measurements by using the Shaeffer's formula (Sastry *et al.*, 1982).

Samples of various feedstuffs available with farmers were collected and analyzed for proximate principles *viz.* dry matter, crude protein, ether extract, crude fibre, nitrogen-free extract and total ash by the standard methods (AOAC, 1995). Nutritive value in terms of DCP and TDN of feedstuffs was calculated using reported values of digestibility coefficients given in the literature (Morrison, 1961; Sen *et al.*, 1978; Pal *et al.*, 1985). Calcium in feed samples was determined by modified method of Talpatra *et al.* (1940). Phosphorus was estimated by Metavandate

method (AOAC, 1995). The carotene content of feeds was calculated by using reported values (Morrison, 1961; Sen *et al.*, 1978). The average availability of carotene was converted to vitamin A (IU) by using formula i.e. 1 mg β -carotene = 400 IU vitamin A activity (Kearl, 1982). Data of feed intake and nutritive values of feed stuffs were used to calculate nutrient (DCP, TDN, Ca, P and Carotene) availability. Then, nutrient availability figures were compared with standard requirements (Paul and Lal, 2010) to work out nutrient excess/deficit.

RESULTS AND DISCUSSION

Survey of existing feeding practices in lactating buffaloes showed that out of 104 dairy farmers visited having 347 buffaloes. All farmers practice stall feeding and animals were fed twice in a day. Five types of feeding patterns were observed as presented in Table 1. Feeding pattern I [Mixture of straw {wheat straw (WS) + gram straw (GS)} + local grasses (LG) + cotton seed cake (CSC) + wheat bran (WB)], Feeding pattern II [wheat straw (WS) + local grasses (LG) + cotton seed cake (CSC) + compounded feed (CF)], Feeding pattern III [Mixture of straw {wheat straw (WS) + soybean straw (SS)} + MP chari + cotton seed cake (CSC) + wheat bran (WB)], Feeding pattern IV [Mixture of straw {wheat straw (WS) + masoor straw (MS)} + maize fodder (MF) + cotton seed cake (CSC) + compounded feed (CF)], and Feeding pattern V [Mixture of straw {gram straw (GS) + masoor straw (MS)} + maize fodder (MF) + cotton seed cake (CSC) + wheat bran (WB)]. The majority of farmers (47%) were feeding mixture of straws (WS + GS) + LG + CSC + WB, followed by WS + LG + CSC + CF (17%), mixture of straws (WS

+ SS) + MP chari + CSC + WB (15%), mixture of straws (WS + MS) + MF + CSC + CF (13%) and mixture of straws (GS + MS) + MF + CSC + WB (8%). Similarly, in villages of Punjab, Haryana and Madhya pradesh wheat straw and cotton seed cake are the predominant feeds for lactating buffaloes, in addition this farmers also use grains and green fodder (Lal *et al.*, 1995; Tomar and Thakur, 2002; Sihag *et al.*, 2002; Mudgal *et al.*, 2003).

The average chemical composition and nutritive value (CP, EE, CF, NFE, TA, AIA Ca, P, DCP and TDN) of available feed stuffs are presented in the Table 2. Results indicated that among dry roughages (wheat, gram, soybean and masoor straws) CP ranged between 3.92 to 6.52%, EE from 0.63 to 1.50%, CF from 33.03 to 41.77%, TA from 6.59 to 11.64%, AIA from 0.69 to 5.84%, Ca from 0.23 to 1.54%, P from 0.04 to 0.15%, DCP from 0.31 to 3.84% and TDN from 42.07 to 53.51%. Among green roughages (local grasses, maize fodder, MP chari) CP varied from 4.29 to 7.00%, EE from 1.14 to 1.48%, TA from 7.95 to 10.69%, AIA from 1.61 to 3.96%, Ca from 0.36 to 0.75%, P from 0.18 to 0.28%, DCP from 1.32 to 4.27% and TDN from 51.15 to 65.87%. Among concentrate feeds (cotton seed cake, wheat bran and compounded feeds) CP ranged between 13.99 to 22.60%, EE 3.45 to 10.22%, CF 9.99 to 27.15%, TA 4.25 to 13.98% and AIA 0.18 to 6.08%. The proximate composition of feedstuffs available for feeding buffaloes in the area of study is comparable with values reported by others (Morrison, 1961; Sen *et al.*, 1978; Kearl, 1982; NRC, 1989; Ranjhan, 1991; NRC 2001; Mudgal *et al.*, 2003; Das *et al.*, 2005; Singh *et al.*, 2005; Anonymous, 2007; Tiwary *et al.*, 2007; Patil *et al.*, 2014; Thakur *et al.*, 2016). The Ca content was similar in available straws and concentrate feeds, except it was much higher in gram straw compared to values reported

by the above workers. The P content in straws, concentrate feeds and green roughages were comparable with values reported by the above workers except it was lower in masoor straw and cotton seed cake than the reported values. Ca: P ratios in rations of buffaloes under different feeding pattern ranged between 1.25:1 to 3.55:1 against optimum ratio 1.56:1 as shown in Table 3. Wider Ca: P ratios in pattern IV (2.76:1) and V (3.55:1) are because of greater proportion of leguminous straw in the rations having higher level of Ca.

Daily requirements and availability of nutrients in different feeding patterns of lactating buffaloes are presented in Table 3. Data indicated excess of DCP (24.11%) and shortage of TDN (4.87%), Ca (4.16%), P (18.17%) and Carotene (18.27%) in feeding pattern I. Similar trends were also observed in feeding pattern II and III i.e. excess of DCP (4.91%) and shortage of TDN (10.93%), Ca (54.83%), P (43.73%) and carotene (17.90%) in feeding pattern II and excess of DCP (10.74%) and shortage of TDN (4.55%), Ca (17.16%), P (20.11%) and carotene (45.11%) in feeding pattern III. While, in feeding pattern IV and V trends were slightly different i.e. excess of DCP (27.87%) and Ca (8.03%) and shortage of TDN (5.24%), P (39.03%) and carotene (57.54%) in feeding pattern IV and similar, excess of DCP (28.52%) and Ca (65.66%) and shortage of TDN (3.28%), P (27.19%) and carotene (51.29%) were also observed in feeding pattern V. In different feeding patterns followed by the farmers for feeding lactating buffaloes in this region, rations of buffaloes was marginally (4.91%) to moderately (28.52%) excess in DCP. Similar excess of DCP was also recorded by earlier workers (Garg *et al.*, 2009; Garg *et al.*, 2013a; Garg *et al.*, 2013b; Garg *et al.*, 2014b; Sherasia *et al.*, 2016) in lactating buffaloes of western and southern region (Gujarat)

of the country. Rations were marginally deficient in TDN (3.28 to 10.93%) and the average daily TDN intake in lactating buffaloes was 7.73 ± 0.02 kg, which was slightly lower than recommended by Paul and Lal (2010). Dietary deficiency of TDN was also observed by others (Lal *et al.*, 1995; Garg *et al.*, 2012; Sherasia *et al.*, 2014). Whereas, Garg *et al.*, 2013a, 2013b; Garg *et al.* (2014b); Sherasia *et al.* (2016) found excess TDN in the ration of lactating buffaloes, while, adequate TDN levels were observed by Kannan *et al.* (2010, 2011) in the ration of lactating buffaloes of Raebareli district in Uttar Pradesh and chitoor district of Andhra Pradesh. Rations of lactating buffaloes were deficient in P (18.17 to 43.73%) and Ca supply was variable. It was deficient in feeding pattern I, II and III and adequate in pattern IV while excess in pattern V depending upon the availability of cereal or leguminous straw. Ca content in leguminous straw was quite higher than cereal straw in the present study. Garg *et al.* (2013b) reported excess Ca. While, others (Garg *et al.*, 2009; Kannan *et al.*, 2010, 2011; Garg *et al.*, 2012; Garg *et al.*, 2013a; Garg *et al.*, 2014ab; Sherasia *et al.*, 2014; Sherasia *et al.*, 2016) were reported shortage of both Ca and P in the ration of lactating buffaloes. There was shortage of carotene by 17.29 to 57.54% in different patterns, due to limited availability of greens. Patil *et al.* (2016); Thakur *et al.* (2016) were also reported deficiency of carotene in the ration of buffaloes of Malwa region of M.P.

CONCLUSION

It was concluded that lactating buffaloes were fed on straw based rations, mainly mixture of cereal (wheat and leguminous straws (gram/masoor/soybean) with little green either local grasses/

Table 1. Feeding patterns and milk yield of lactating buffaloes.

S.No.	Name of village	No. of farmers	No. of buffaloes	Avg. Milk yield (kg/h/d)	Feeding patterns and no. of farmers				
					(WS+GS)+ local grasses+CSC+WB (I)	WS+local grasses+CSC+CF (II)	MP chari+CSC+WB (III)	(WS+MS)+Maize fodder+CSC+CF (IV)	(GS+MS)+Maize fodder+CSC+WB (V)
1	Bhaislai	11	33	7.52	6	2	2	1	-
2	Sonwai	10	25	6.80	8	1	1	-	-
3	Borkhedi	10	30	7.17	6	2	-	1	1
4	Panda	12	40	7.50	2	3	1	4	2
5	Navda	9	32	7.80	4	2	2	1	-
6	Harsola	12	42	7.60	5	1	2	2	2
7	Santer	11	43	7.91	4	2	3	1	1
8	Kavti	12	55	7.27	6	3	2	-	1
9	Rangwasa	09	23	8.50	5	1	2	1	-
10	Umriya	08	24	6.36	3	1	1	2	1
	Total	104	347	-	49	18	16	13	8
	%	-	-	-	47.12	17.31	15.38	12.50	7.69

WS = Wheat Straw, GS = Gram Straw, MS = Masoor Straw, SS = Soybean Straw, CSC = Cotton Seed Cake, CF = Compounded Feed

Table 2. Average chemical composition and nutritive value of feed stuffs (% DM basis).

Feed stuffs	CP	EE	CF	NFE	TA	AIA	Ca	P	Carotene (ppm)	DCP	TDN
Wheat straw	3.92±0.22	0.99±0.06	33.07±0.61	50.38±0.66	11.64±0.64	5.84±0.16	0.23±0.02	0.06±0.30	0.20	0.31±0.01	42.07±0.27
Gram straw	6.24±0.24	0.63±0.50	39.16±0.74	42.95±1.20	7.98±0.34	2.44±0.15	1.54±0.08	0.04±0.01	1.00	2.80±0.11	42.81±0.27
Masoor straw	6.52±0.32	1.50±0.11	36.91±0.82	46.21±0.87	8.84±0.30	4.17±0.23	1.46±0.06	0.05±0.01	1.00	3.84±0.19	53.51±0.23
Soybean straw	6.14±0.34	0.80±0.06	41.77±1.46	45.52±1.74	6.59±0.38	0.69±0.06	0.94±0.04	0.15±0.02	Traces	1.77±1.10	43.76±0.55
Local grasses (Baroo)	4.29±0.67	1.48±0.04	26.94±0.45	56.74±0.60	10.55±0.01	3.61±0.26	0.36±1.02	0.18±0.01	94.48	2.44±0.09	51.15±0.76
Maize fodder (32% DM)	7.00±0.03	1.14±0.04	30.64±0.64	50.55±0.02	10.69±0.76	1.61±0.84	0.75±0.01	0.28±0.01	58.74	4.27±0.02	65.87±0.03
MP chari	6.30±0.02	1.18±0.01	31.62±0.23	52.95±0.11	7.95±0.09	3.96±0.31	0.61±0.20	0.18±0.01	67.71	1.32±0.10	55.84±0.05
Cotton seed cake	22.60±0.42	10.22±0.64	27.15±1.72	35.75±1.15	4.25±0.22	0.18±0.02	0.22±0.01	0.51±0.02	0.20	19.02±0.38	78.96±1.48
Wheat bran	13.99±0.52	3.45±0.11	9.99±0.92	68.06±1.63	4.51±0.74	0.48±0.10	0.21±0.01	0.61±0.05	0.10	11.32±0.42	74.85±0.58
Compounded feed	18.25±0.06	3.41±1.20	12.20±0.97	48.63±0.08	14.77±0.06	2.59±0.56	0.32±0.05	0.73±0.02	NA	13.14±0.12	64.18±0.06

NA=Not available

Table 3. Average daily requirement and availability of nutrients in different feeding patterns for lactating buffaloes.

Feeding pattern I [Mixture of Straw (WS+GS)+LG+CSC+WB]											
	Body weight (Kg)	Milk yield (Kg)	Milk fat (%)	DM (Kg)	DCP (g)	TDN (Kg)	Ca (g)	P (g)	Ca: P Ratio	Carotene (ppm)	
1	2	3	4	5	7	8	9	10	11	12	
Requirement (Paul and Lal, 2010)	575	8.50	6.50	13.19	864.83	7.81	69.91	44.85	1.56:1	109.25	
Availability (Per head/day)	562±8.10	8.50±1.20	6.50±0.80	12.53±0.61	1073.30±4.62	7.43±0.03	67.00±0.21	36.70±0.10	1.83:1	89.29	
Excess (+) /Deficient (-)					(+) 208.47	(-) 0.38	(-) 2.91	(-) 8.15	-	(-) 19.96	
Deficiency (-) /Excess (+) (%)					(+) 24.11	(-) 4.87	(-) 4.16	(-) 18.17	-	(-) 18.27	
Feeding pattern II (WS+LG+CSC+CF)											
Requirement (Paul and Lal, 2010)	550	7.25	6.00	11.74	757.49	6.95	62.22	39.99	1.56:1	104.50	
Availability (Per head/day)	551±7.10	7.13±0.03	5.90±0.10	11.16±0.51	794.70±5.62	6.19±0.01	28.10±0.20	22.50±0.15	1.25:1	85.79	
Excess (+) /Deficient (-)					(+) 37.21	(-) 0.76	(-) 34.12	(-) 17.49	-	(-) 18.71	
Deficiency (-) /Excess (+) (%)					(+) 4.91	(-) 10.93	(-) 54.83	(-) 43.73	-	(-) 17.90	
Feeding pattern III [Mixture of straw (WS+SS)+MP charri+CSC+WB]											
Requirement (Paul and Lal, 2010)	525	6.50	5.50	10.75	682.50	6.37	56.98	36.55	1.56:1	99.75	
Availability (Per head/day)	515±6.50	6.20±0.05	5.45±0.12	10.61±0.62	755.80±6.65	6.08±0.02	47.20±0.24	29.20±0.14	1.62:1	54.66	
1	2	3	4	5	7	8	9	10	11	12	
Excess (+) /Deficient (-)					(+) 73.30	(-) 0.29	(-) 9.78	(-) 7.35	-	(-) 45.09	
Deficiency (-) /Excess (+) (%)					(+) 10.74	(-) 4.55	(-) 17.16	(-) 20.11	-	(-) 45.20	
Feeding pattern IV [Mixture of straw (WS+MS)+MF+CSC+CF]											
Requirement (Paul and Lal, 2010)	550	7.00	6.00	11.58	743.00	6.86	61.37	39.37	1.56:1	104.50	
Availability (Per head/day)	548±6.10	6.73±0.03	5.80±0.14	11.31±0.41	950.10±5.60	6.50±0.04	66.30±0.18	24.00±0.11	2.76:1	44.37	
Excess (+) /Deficient (-)					(+) 207.10	(-) 0.36	(+) 4.93	(-) 15.37	-	(-) 60.13	
Deficiency (-) /Excess (+) (%)					(+) 27.87	(-) 5.24	(+) 8.03	(-) 39.03	-	(-) 57.54	

Table 3. Average daily requirement and availability of nutrients in different feeding patterns for lactating buffaloes. (Continue)

Feeding pattern V [Mixture of straw (GS+MS)+MF+CSC+WB]											
Requirement (Paul and Lal, 2010)	550	6.50	6.00	11.23	716.00	6.70	59.52	38.18	1.56:1	104.50	
Availability (Per head/day)	541±7.20	6.20±0.01	5.80±0.14	10.60±0.41	920.20±6.30	6.48±0.06	98.60±0.18	27.80±0.13	3.55:1	50.90	
Excess (+) /Deficient (-)				(-) 0.63	(+) 204.20	(-) 0.22	(+) 39.08	(-) 10.38	-	(-) 53.60	
Deficiency (-) /Excess (+) (%)					(+) 28.52	(-) 3.28	(+) 65.66	(-) 27.19	-	(-) 51.29	

MP chari/ maize fodder and un-decorticated cotton seed cake/wheat bran/compounded feed as concentrate. DCP was moderately high and TDN was marginally deficient while phosphorous and carotene were moderate to highly deficient in the rations of lactating buffaloes of the region. Calcium was deficient in feeding pattern I, II and III and adequate in pattern IV while excess in pattern V.

REFERENCES

- Anonymous. 2007. Chemical composition of different straws. Available on www.pubmed.com.
- Association of Official Analytical Chemists (AOAC). 1995. *Official Methods of Analysis*, 15th ed. Association of Official Analytical Chemists, Washington, DC, USA.
- Das, M.M., S.B. Maity and S. Kundu. 2005. Evaluation of masoor straw based ration in sheep and goats. *Indian J. Anim. Nutr.*, **22**(3): 204-205.
- Garg, M.R., S.P. Arora, B.M. Bhanderi, P.L. Sherasia and D.K. Singh. 2000. Mineral status of feeds and fodders in Khaira district of Gujarat. *Indian J. Dairy Sci.*, **53**: 291-297.
- Garg, M.R., S.A. Biradar and A. Kannan. 2009. Assessment of economic impact of implementing ration balancing programme in lactating cows and buffaloes under field conditions. *Indian J. Anim. Nutr.*, **26**(2): 146-150.
- Garg, M.R., A. Kannan, B.T. Phondba, S.K. Shelke and P.L. Shearson. 2012. A study on the effect of ration balancing for improving milk production and reducing methane emission in lactating buffaloes under field conditions. *Indian J. Dairy Sci.*, **65**(3): 250-255.
- Garg, M.R., P.L. Sherasia, B.M. Bhanderi and B.T. Phondba. 2014a. Effect of ration balancing on methane emission, faecal Archaeol concentration and its relation to enteric methane in crossbred cows. *Indian J. Anim. Sci.*, **84**(6): 687-690.
- Garg, M.R., P.L. Sherasia, B.M. Bhanderi, B.T. Phondba, S.K. Shelke and H.P.S. Makkar. 2013a. Effects of feeding nutritionally balanced rations on animal productivity, feed conversion efficiency, feed nitrogen use efficiency, rumen microbial protein supply, parasitic load, immunity and enteric methane emissions of milking animals under field conditions. *Anim. Feed Sci. Tech.*, **179**: 24-35.
- Garg, M.R., P.L. Sherasia, B.M. Bhanderi, B.T. Phondba, S.K. Shelke and C.T. Patel. 2013b. Effect of feeding balanced rations on milk production, enteric methane emission and metabolic profile in crossbred cows under field conditions. *Indian J. Dairy Sci.*, **66**(2): 113-119.
- Garg, M.R., P.L. Sherasia, B.T. Phondba and S.A. Hossain. 2014b. Effect of feeding a balance ration on milk production, microbial nitrogen supply and methane emission in field animals. *Anim. Prod. Sci.*, **54**: 1657-1661.
- Kannan, A., M.R. Garg and B.V.M. Kumar. 2011. Effect of ration balancing on milk production, microbial protein synthesis and methane emission in crossbred cows under field conditions in Chittoor district of Andhra Pradesh. *Indian J. Anim. Nutr.*, **28**(2): 117-123.
- Kannan, A., M.R. Garg and P. Singh, 2010. Effect of ration balancing on methane emission and milk production in lactating animals

- under field conditions in Rae Bareilly district of Uttar Pradesh. *Indian J. Anim. Nutr.*, **27**(2): 103-108.
- Kearl, L.C. 1982. *Nutrient Requirements of Ruminants in Developing Countries*. International Feed Stuffs Institute, Utah Agriculture Experimental Station. Utah State University, Logan, Utah, USA.
- Lal, D., R. Gupta, V.B. Dixit and R.K. Sethi. 1995. Nutritional status of lactating buffaloes in Hisar-A random survey analysis. *Indian J. Dairy Sci.*, **48**: 378.
- Morrison, F.B. 1961. *Feed and Feeding, Abridged*, 9th ed. The Morrison Publishing Company, Canada.
- Mudgal, V., M.K. Mehta, A.S. Rane and S. Nanawati. 2003. A survey on feeding practices and nutritional status of dairy animals in Madhya Pradesh. *Indian J. Anim. Nutr.*, **20**(2): 217-220.
- NRC. 1989. *Nutrient Requirements of Dairy Cattle*, 6th ed. National Academic of Sciences. National Research Council, Washington. D.C., USA.
- NRC. 2001. *Nutrient Requirements of Dairy Cattle*, 7th ed. National Academic of Sciences. National Research Council, Washington. D.C., USA.
- Pal, S., C.S. Rathi and S.M. Chahal. 1985. Chemical composition of masoor straw. *Indian J. Anim. Sci.*, **55**: 832.
- Patil, N., R.K. Jain and V. Mudgal. 2014. Effect of strategic nutrient supplementation on the reproductive performance of anoestrus buffaloes in the Malwa region of Madhya Pradesh. *Buffalo Bull.*, **33**(2): 199-207.
- Patil, N., R.K. Jain and D. Tewari. 2016. Nutritional status and hemato-biochemical profile of anoestrus buffaloes of Malwa region of Madhya Pradesh. *Buffalo Bull.*, **35**(3): 417-426.
- Paul, S.S. and D. Lal. 2010. *Nutrient Requirements of Buffaloes*. Satish Serial Publishing House, New Delhi, India. 137p.
- Ranjhan, S.K. 1991. *Chemical Composition and Nutritive Value of Indian Feeds and Feeding of Farm Animals*. Indian Council of Agricultural Research Publication, New Delhi, India.
- Sastry, N.S.R., C.K. Thomas and R.A. Singh. 1982. *Farm Animal Management and Poultry Production*, 2nd ed. Vikas Publishing House Pvt Ltd. New Delhi, India.
- Sen, K.C., S.N. Ray and S.K. Ranjhan. 1978. *Nutritive Value of Indian Feeds and Feeding of Farm Animals*. Indian Council of Agricultural Research, New Delhi, India.
- Sherasia, P.L., M.R. Garg and B.M. Bhandari. 2014. Effect of feeding balanced rations on production efficiency and enteric methane emission in lactating buffaloes under tropical conditions. *Buffalo Bull.*, **33**(4): 423-431.
- Sherasia, P.L., B.T. Phondba, S.A. Hossain and M.R. Garg. 2016. Effect of feeding balanced rations on milk production, methane emission metabolite and feed conversion efficiency in lactating cow. *Indian J. Anim. Res.*, **50**(4): 505-511.
- Sihag, S., K.R. Yadav, S.S. Khirwar, Rajesh and H. Lal. 2002. Availability of feeds and feeding practices and socio-economic status of farmers in irrigated cotton based system. *Indian J. Anim. Nutr.*, **19**(3): 256-261.
- Singh, B., J.L. Chaudhary and N.K. Rajora. 2005. Nutritive evaluation of soybean straw in sheep and goats. *Indian J. Anim. Nutr.*, **22**(1): 67-68.

- Talpatra, S.K., S.C. Ray and K.C. Sen. 1940. Analysis of mineral constituents in biological materials. *Indian Journal of Veterinary Science and Animal Husbandry*, **10**: 243-245.
- Tewari, D., R.K. Jain and N. Patil. 2012. Feeding pattern, prevalence of reproductive disorders and nutritional status of available feed stuffs for dairy animals in Indore district of M.P. *Indian J. Anim. Nutr.*, **4**: 356-359.
- Tewari, D., R.K. Jain and V. Mudgal. 2014. Effect of strategic nutrient supplementation on the reproductive performance of anoestrous crossbred cattle in Malwa region of Madhya Pradesh. *Indian J. Anim. Res.*, **48**(6): 580-584.
- Tiwary, M.K., D.P. Tiwari, A. Kumar and B.C. Mondal. 2007. Existing feeding practices, nutrient availability and reproductive status of dairy cattle and buffaloes in Haridwar district of Uttarakhand. *Indian J. Anim. Nutr.*, **7**: 177-185.
- Tomar, S.K. and S.S. Thakur. 2002. Feed resources, feeding practices, milk production and disposal pattern in Karnal district. *Indian J. Dairy Sci.*, **55**: 306-309.